
THE IMPORTANCE OF FOOD ITEM TO MAKE AN ENERGETIC ATHLETES

DR.B.ASHOK KUMAR

ASSISTANT PROFESSOR

DEPARTMENT OF PHYSICAL EDUCATION

AND SPORTS SCIENCES

ANNAMALAI UNIVERSITY

Abstract

Dairy products are thought to improve recovery after both resistance and endurance exercises due to their nutritional properties. We systematically reviewed the effects of dairy product intake on exercise performance and recovery of muscle function in humans. A literature search was conducted in the MEDLINE (via PubMed) and Web of Science databases from their inception to 15th April 2018. The initial search retrieved 7708 articles, and a total of 11 studies were finally included after applying inclusion and exclusion criteria. All the selected studies were conducted with cow's milk. Whereas some studies found significant positive effect of cow's milk on exercise performance and recovery of muscle function, others did not find any effect. These controversies could be due to the heterogeneity of cow's milk ingestion (e.g., amount of cow's milk, timing of consuming the cow's milk), to the type of intervention, and to the large heterogeneity of outcomes measured. Limited studies exist examining the effects of cow's milk consumption and its influence on exercise performance and recovery of muscle function, therefore further studies are needed to draw more definitive conclusions.

Keywords: Dairy product, Muscle damage, Muscle recovery, Resistance training, Endurance training

Introduction

There is evidence that proper nutritional intake is a key factor in optimizing exercise performance as well as adaptation to training (e.g., positive stimuli for protein synthesis in skeletal muscle) and recovery of muscle function (e.g. increase the recovery between training sessions or competitions, decrease the symptoms of delayed onset muscle soreness, etc.) [1, 2]. High exercise performance requires very controlled nutritional intake [3] and timing [4] before, during and after exercise to maximize exercise-induced adaptation and to shorten recovery after exercise, however the impact of either the type, composition or timing of the nutrient is still not known. Protein intake has a great impact on muscle damage repair, facilitating the recovery of muscle function (e.g. muscle strength, muscular power production, muscular stiffness, etc.) and muscular protein synthesis [1, 2]. For both hypertrophy and recovery, a positive muscle protein net balance, i.e., a higher muscle protein synthesis than muscle protein breakdown, is necessary [5].

When the rates of muscle protein synthesis and degradation increase [6, 7], an adequate nutrition is required [8– 10] to facilitate the recovery process. For example, a bout of unaccustomed exercise, especially that including eccentric muscle contractions such as downhill running, can damage contractile proteins, impair muscle function, and induce muscle soreness [11, 12]. In theory, the stimulation of muscle protein synthesis by means of protein or amino acids (e.g. through dairy products ingestion) represents an important skeletal muscle adaptive response to mechanical stress that helps in recovery of muscle function [8, 13, 14].

Dairy products are rich in amino acids, proteins, lipids, minerals and vitamins, and their health benefits have been reviewed elsewhere [15]. These beneficial properties are based on the fact that dairy products, and especially cow's

1“PROmoting Fitness and Health through physical activity” research PROFITH group
Department of Physical Education and Sport, Faculty of Sport

Sciences, University of Granada, Ctra. de Alfacar s/n C.P, 18071 Granada, Spain milk, contains lactose (carbohydrate), casein and whey protein—commonly in a 3:1 ratio (casein:

whey), as well as calcium [5, 15]. Of note is that these other nutrients present in cow's milk such as calcium, sodium or potassium could

Full list of author information is available at the end of the article

The Author(s). 2019 Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Aid in fluid recovery after exercising [5] and this improvement in the hydration state could help the recovery of the skeletal muscle. Furthermore, the aforementioned protein ratio could promote slow digestion and absorption of amino acids [5], which may lead to an increase in the serum amino acid concentration (mainly branched amino acids) [16], however, it is important to note that casein alone or whey protein alone, could increase serum amino acid concentration. These branched amino acids may have a large impact on protein synthesis and muscle metabolism [5] and therefore, helping the aforementioned muscle damage repair process. However, it is important to note that Atherton et al. [17] showed that branched amino acids effect on muscular protein synthesis is most likely due to the presence of leucine and not the presence of isoleucine or valine. Furthermore Witard et al. [18] reported that muscular protein synthesis stimulation via branched amino acids was ~ 50% inferior compared to a whey protein bolus containing similar amounts of branched amino acids. Moreover, the nutritional characteristics of dairy products (e.g. cow's milk) [19] plus the relatively low price and high availability [20] of dairy products make them a potentially recovery-enhancing product after exercise [5]. This is observed in the current growth of scientific interest in the effects of dairy product intake on exercise performance and muscle function recovery [5].

In this systematically review, we summarize the results of the studies assessing the effect of dairy products on exercise performance and on the recovery of muscle function in humans.

Methods

This systematic review was conducted following the Preferred Reporting Items for Systematic Reviews and MetaAnalysis (PRISMA) statement [21] and was registered through the International Prospective Register of Systematic Reviews (PROSPERO registration number: CRD4201 8094800).

Search strategy

A literature search was conducted in the MEDLINE (via PubMed) and Web of Science (WOS) databases from their inception to 15th April 2018. The search terms as well as the search strategy and equations can be seen in detail in the Additional file 1: Table S1. Briefly, we used “dairy products”, “exercise”, “training”, “athletic performance”, “muscle strength”, “muscle fatigue”, and “muscle recovery” among others terms (see Additional file 1: Table S1) joined with Boolean operators. The reference lists of the retrieved systematic reviews and meta-analyses were reviewed to identify additional studies.

Selection criteria

The inclusion criteria used were 1) dairy product and exercise intervention (either chronic or acute) studies. The difference between the intervention and the control group/ period should be in the dairy product consumption. Dairy product includes raw and processed or manufactured milk and milk-derived products. Dairy products normally come from cow but could be also from goats, sheep, reindeer, and water buffalo as defined by the National Library of Medicine (PubMed) [22]; 2) conducted in healthy humans, regardless of age or fitness level; and 3) studies including measurements of exercise performance or recovery of muscle function. We included studies that measured exercise performance quantified by fitness parameters such as maximum repetition test and isokinetic dynamometry variables [23, 24]. Moreover, we included studies that assessed muscle recovery function by subjectively measurements [e.g., ratio of perceived exertion and visual analogue scales (VAS)] or objectively measured by the use of blood markers [(e.g., creatine kinase (CK) and

myoglobin) [25]. If the same data/study was used in different original articles for different purposes, only the report that provided more detailed information about the topic of this systematic review was included.

The exclusion criteria used were 1) studies written in languages other than English or Spanish; 2) studies in which any type of protein, flavoring or sweetener was added to the consumed dairy product; moreover, colostrum (e.g., bovine colostrum), chocolate milk and breast milk were excluded from this systematic review; 3) studies in which there was no a control group.

Data extraction

The following data were collected from each included study: 1) study characteristics (author identification and reference); 2) number of participants and sex; 3) age of the participants; 4) fitness level of the participants; 5) design; 6) groups; 7) exercise intervention; 8) dairy product ingestion (e.g. cow's milk ingestion) and placebo ingestion; 9) study outcomes; 10) results; and 11) risk of bias score.

Regarding the exercise intervention, those studies including exercises such as sprints series, isokinetic (combining eccentric and concentric contractions) or resistance (e.g. bench press) exercise or training were classified in resistance or high-intensity exercise. Those studies including exercises such as continuous cycling or cycling at different intensities (e.g. 70% peak oxygen uptake) were classified in endurance exercise.

Study quality and risk of bias assessment

The Cochrane risk of bias tool [26] was used to evaluate the risk of bias in each study. This tool assesses random sequence generations and allocation concealment, performance bias (blinding of participants and personnel), detection bias (blinding of the outcome assessment), attrition bias (incomplete outcome data), reporting bias (selective reporting), and other sources of bias.

The literature search and data extraction as well as the quality assessment were independently performed by four reviewers (JMAA, GSD, BMT and JRR), and inconsistencies were solved by consensus.

Results

Overall results

Figure 1 presents the PRISMA consort diagram for the search strategy. The initial search retrieved 7708 articles, and a total of 11 studies were finally included after applying the inclusion and exclusion criteria (Table 1). All the included studies used cow's milk products. A total of 7 studies (63.6%) investigated the acute effect of cow's milk after resistance or high-intensity exercise [27–33], whereas three studies (27.3%) determined the acute effect of cow's milk after endurance exercise [34–36]. In addition, one study [37] analyzed the long-term effect (over 12 weeks of resistance training) of exercise and cow's milk on the maximum repetition strength of squat and bench press.

Risk of bias within studies

The quality of the included studies was predominantly suboptimal. The methodological quality assessment is shown in Fig. 2. Details of randomization [27–37], allocation concealment [27–37], and study blinding [27–32, 35–37] were inadequately reported or rated as an “unclear risk” (categorized when the information was not specified in the article) for most studies.

Effects of cow's milk on performance and muscle function recovery after resistance or high-intensity exercise

The exercise performance outcomes measured were muscular strength related variables [27–33, 37]. Regarding muscle function recovery, most of the studies [27–33] determined muscle soreness or damage using subjective scales such as VAS, and the most of them added also measures of blood biomarkers (e.g., CK or myoglobin) [27–32].

Cow's milk attenuated losses in peak torque (maximal effort concentric knee flexion) [27, 28, 30, 31], total work of the set (6 concentric knee flexion repetitions) using isokinetic

dynamometry [31], countermovement jump [27, 28, 33], rate of force development of an isometric contraction of the dominant leg quadriceps [27, 28], and sprint tests [27, 29, 32]. On the other hand, other studies observed no effect of cow's milk on sprint recovery [28, 33], countermovement jump [32], reactive strength index [32], and bench and squat maximum strength after an exercise intervention [37].

Cow's milk did not modify the pre-post resistance exercise changes in serum CK [27, 28, 32], myoglobin [32], high-sensitivity C-reactive protein [27, 28] and protein carbonyls [27]. In contrast, a positive effect, i.e. lower increase in CK and myoglobin concentrations, was observed from the baseline to 48 h after resistance exercise in the cow's milk group compared with the values of the placebo beverage group [31]. Cockburn et al. [30] also showed that the increase in CK can be blunted after resistance exercise with less cow's milk ingestion (500 mL of cow's milk instead of 1000 mL of cow's milk). These lower increases of CK were observed from 24 to 72 h after exercise-induced muscle damage in the hamstring and cow's milk ingestion [29]. Cow's milk also attenuated the skeletal troponin I increase after exercise compared with a placebo group (energy-matched carbohydrate solution) [29].

Cow's milk did not improve muscle soreness after resistance exercise in other studies [31–33], whereas it had a positive effect on muscle soreness and tiredness at 72 h post resistance exercise in other [27]. Moreover, cow's milk reduced passive soreness in males and females, as well as active muscle soreness (all from baseline to 72 h) in both sexes [29]. Similar results were found in another study that compared cow's milk vs. energy-matched carbohydrate solution as a control at 72 h [28]. Finally, passive measurements of muscle soreness (using VAS) showed a benefit of limiting increases in muscle soreness in the group receiving less bolus cow's milk (500 mL) compared with the high-bolus cow's milk group (1000 mL) between the baseline and 48 h after exercise and cow's milk ingestion [30]. Due to lack of homogeneity in the measurement of exercise performance, and on the recovery of muscle function outcomes after resistance or high-intensity exercise intervention doing a meta-analysis was not possible (see Table 1).

Effects of cow's milk on exercise performance and muscle function recovery after endurance exercise Cow's milk before (2 h prior the exercise) endurance exercise improved

performance in a 20-km time trial ($P < 0.05$) [34]. Moreover, no differences in heart rate were observed between the cow's milk group and the placebo group [34]. In another study [35], the mean cycling time to exhaustion was the same on the placebo group trial (39.6 ± 7.3 min) compared with the cow's milk group (39.7 ± 8.1 min; $P = 0.879$). Furthermore, no differences in oxygen consumption during exercise were found [35]. Finally, there was no effect of cow's milk on changes in the rate of perceived exertion after exercise ($P = 0.744$) compared with the placebo group [35].

Lee et al. [36] did not observe differences in the time to volitional exhaustion independently of the beverage ingested (median, range: 103.3, 85.7–228.5 vs. 93.3, 82.4– 192.3 min for the cow's milk vs. placebo group, respectively) [36]. Cow's milk did not alter the heart rate reached in the volitional exhaustion point nor the RPE sensations during the exercise [36]. Due to heterogeneity in the measurement of exercise performance and on the recovery of muscle function outcomes after endurance exercise doing a meta-analysis was not possible (see Table

Discussion

We systematically reviewed and summarized the results of the studies investigating the effects of dairy products on exercise performance or on the recovery of muscle function in humans. All the studies meeting the criteria were conducted using cow's milk. The studies investigating the effect of cow's milk after high-intensity exercise [27–33] or resistance training [37] reported contradictory results. Whereas some studies found significant effects of cow's milk intake on performance or recovery of muscle function such as attenuated losses in peak torque, countermovement jump, rate of force development, sprint series and inflammation and markers of muscle damage, others did not find any effect. Therefore, there is currently not enough evidence to conclude whether cow's milk has a positive effect on exercise performance and recovery of muscle function, and further studies are needed to make more definitive conclusions. The results regarding endurance exercise are similar to those observed in resistance exercise. Whereas one study observed significant effect of cow's milk intake on performance [34] others [35, 36] did not observe any enhancing effect. The observed contradictory findings can be explained by the heterogeneity of cow's milk

ingestion, such as the amount of cow's milk, the timing of the cow's milk intake, and the type of intervention, as well as by the large heterogeneity of outcomes measured. Moreover, the study participants' fitness level may also play an important role when studies are compared. In addition, doing a meta-analysis was not appropriate due to the heterogeneity between studies, which was mainly identified in the reporting of exercise performance and on the recovery of muscle function.

It has been shown that 20 g of protein might be sufficient to stimulate muscle protein synthesis after resistance exercise [38], so perhaps a greater consumption of protein (via cow's milk) would have resulted in more positive effects. It is noteworthy, however, that one study showed similar effects after consumption of 500 and 1000 mL of cow's milk [30]. Volek et al. [37] found no differences between the effects of cow's milk vs. juice (both groups consumed 708 mL daily) after a resistance exercise intervention in maximum repetition strength (12 weeks of resistance training). Both Lee et al. [36] and Watson et al. [35] found no effect of cow's milk on time to volitional exhaustion, heart rate, expired gases and RPE after endurance exercise. In contrast, Upshaw et al. [34] found that low-fat cow's milk (2262 ± 299 mL) improved the time in a 20-km time trial test after glycogen-lowering exercise compared with that of a placebo group (2262 ± 290 mL low-energy drink). Regarding the intensity of the exercise, one study [35] reported that the heart rate during endurance exercise in a cow's milk trial test (2263 ± 241 mL) was higher than that during a carbohydrate trial test (2280 ± 249 mL), yet no differences in oxygen consumption during exercise were observed. The intensity and exercise performed in both studies was similar [35, 36], while in the study by Upshaw et al. [34], a glycogen-lowering exercise was performed before both cow's milk ingestion and the examined exercise. The amount of cow's milk ingested in the study by Upshaw et al. [34] and in the study by Watson et al. [35] was similar, and therefore, the differences in recovery between the groups might be partially explained by differences in the fitness levels of the participants [35, 36]. In the study by Upshaw et al. [34], the participants were trained cyclists, while in the other study, the participants were regularly active individuals.

Philips et al. [39] in their review focused on the evidence showing the differences in responses of muscle protein synthesis and muscle protein accretion in humans, concluded

that cow's milk-based proteins (whey and casein) appear to be better than carbohydrate beverages in the promotion of hypertrophy. Moreover, they highlighted the importance of the dose-response in the studies since the difference in leucine content (as is present in cow's milk) may have an important influence in maintaining and possibly increasing the muscle mass [39]. Furthermore, leucine could have an impact in the recovery process (e.g. in the muscle protein synthesis and muscle protein accretion). Maybe the contradictory results obtained in exercise performance in our systematic review could be partially explained by the different amount of cow's milk (and therefore leucine content) provided in the selected studies. It is assumed that a single cow's milk bolus intake increases net amino acid synthesis in young healthy sedentary volunteers [40]. Moreover, the consumption of protein and carbohydrates together, as is presented in cow's milk, results in a higher rate of protein synthesis compared to that from the intake of these nutrients separately [41, 42]. Regarding protein consumption, sufficient protein intake is necessary to stimulate protein synthesis as we mentioned previously [38]. Therefore, it is biologically plausible that cow's milk consumption after a bout of exercise can stimulate protein synthesis metabolism [6, 7]. Muscle membrane damage following exercise occurs as a result of mechanical stress during the first phase of muscular damage, with further disruption via the lysosomal pathway during the next phase [43]. The lack of a positive effect of cow's milk on some blood markers (e.g., protein carbonyls and high-sensitivity C-reactive protein) [27, 28, 32] may suggest that the ingestion of cow's milk could affect other metabolic pathways. For example, the CK concentration increases after exercise, but no clear CK blunting effects were observed from cow's milk, while the skeletal troponin I rise was blunted in the cow's milk group [29]. Controversially, in another study it was shown that the CK increase can be attenuated with lower cow's milk intake (500 mL instead of 1000 mL) [30], and an effect on myoglobin was also observed between the baseline and 48 h after exercise [31] from cow's milk ingestion. Of note is that in both of these studies, participants played regularly in team sports, and this fact could result in less muscle damage due their fitness level. In their review, Sousa et al. [44] recommend the ingestion of 0.8–1.2 g carbohydrate/kg/h and 0.2–0.4 g protein/kg/h preferably after the exercise, with a minimum of 20 g high-quality protein for improve the recovery after exercise. However, some controversial still exist regarding the correct timing and if carbohydrate and protein have to be consumed right after the exercise [44, 45].

Following these recommendations by Sousa et al. [44], maybe the negative results found in some of the included studies could be partially explained for either an insufficient amount of cow's milk (i.e. not enough protein and/or leucine) or the timing of ingestion. Regarding the time of ingestion, most of the studies provided the beverage (e.g. cow's milk, placebo, etc.) immediately after the exercise [27– 32], while others offered the beverage during the exercise [35, 36], and one [34] offered the beverage immediately after the exercise and every 30 min. In Volek et al. [37] the beverage consumption was daily.

Regarding muscle soreness perception, whereas several studies did not find a positive effect of cow's milk [31–33, 36], others observed significant differences in both active and passive muscle soreness between cow's milk and control groups (~ 500 mL) [27–30]. However, VAS muscle soreness perception is a more subjective outcome than blood markers, and it is more difficult to establish whether these differences could be explained by the treatment (e.g., cow's milk vs. placebo), by the physical condition of participants (e.g., team sport players vs. individuals not habituated to exercise), or by unaccustomed sensations after resistance (i.e., repeated eccentrics contractions).

Conclusions

In conclusion, based on the current evidence, it cannot be determined whether cow's milk has a positive effect on exercise performance and recovery of muscle function in humans, due to the limited number of studies included in this systematic review. Nevertheless, since cow's milk is a source of protein, carbohydrates, calcium and other nutrients, and thus may lead to an increase in the serum amino acid concentration and, therefore, helping the muscle damage repair process. In line with this, some studies included found significant effects of cow's milk intake on performance and recovery of muscle function. For these reasons, more and better study designs such as blinding the beverage to both, participants and personnel, generate a random sequence of beverage group, etc. are needed to demonstrate its usefulness as a sport nutrition-related supplement.

References

1. Pennings B, Koopman R, Beelen M, Senden JMG, Saris WHM, van Loon LJC. Exercising before protein intake allows for greater use of dietary protein-derived amino acids for de novo muscle protein synthesis in both young and elderly men. *Am J Clin Nutr.* 2011;93:322–31.
2. Yang Y, Breen L, Burd NA, Hector AJ, Churchward-Venne TA, Josse AR, et al. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr.* 2012;108:1780–8.
3. Ziegenfuss TN, Landis JA, Lemieux RA. Protein for sports-new data and new recommendations. *Strength Cond J.* 2010;32:65–70.
4. Areta JL, Burke LM, Ross ML, Camera DM, West DWD, Broad EM, et al. Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis. *J Physiol.* 2013; 591:2319–31 Wiley-Blackwell.
5. Roy BD. Milk: the new sports drink? A review. *J Int Soc Sports Nutr.* 2008;5:15.
6. Pitkanen HT, Nykanen T, Knuutinen J, Lahti K, Keinanen O, Alen M, Komi PV, Mero AA. Free amino acid pool and muscle protein balance after resistance exercise. *Med Sci Sports Exerc.* 2003;35:784–92.
7. Biolo G, Maggi SP, Williams BD, Tipton KD, Wolfe RR. Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol Metab.* 1995;268:E514–20.
8. Levenhagen DK, Carr C, Carlson MG, Maron DJ, Borel MJ, Flakoll PJ. Postexercise protein intake enhances whole-body and leg protein accretion in humans. *Med Sci Sports Exerc.* 2002;34:828–37.
9. Koopman R, Wagenmakers AJM, Manders RJF, Zorenc AHG, Senden JMG, Gorselink M, et al. Combined ingestion of protein and free leucine with carbohydrate

increases postexercise muscle protein synthesis *in vivo* in male subjects. *Am J Physiol Metab.* 2005;288:E645–53.

10. Rasmussen BB, Tipton KD, Miller SL, Wolf SE, Wolfe RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol.* 2000;88:386–92.
11. Clarkson PM, Nosaka K, Braun B. Muscle function after exercise-induced muscle damage and rapid adaptation. *Med Sci Sports Exerc.* 1992;24:512–20.
12. Pasiakos SM, Lieberman HR, McLellan TM. Effects of protein supplements on muscle damage, soreness and recovery of muscle function and physical performance: a systematic review. *Sports Med.* 2014;44:655–70.
13. Moore DR, Stellingwerff T. Protein ingestion after endurance exercise: the “evolving” needs of the mitochondria? *J Physiol.* 2012;590:1785–6 Wiley-Blackwell.
14. Howarth KR, Moreau NA, Phillips SM, Gibala MJ. Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans. *J Appl Physiol.* 2009;106:1394–402.
15. Haug A, Høstmark AT, Harstad OM. Bovine milk in human nutrition – a review. *Lipids Health Dis.* 2007;6:25.
16. Bos C, Metges CC, Gaudichon C, Petzke KJ, Pueyo ME, Morens C, et al. Postprandial kinetics of dietary amino acids are the Main determinant of their metabolism after soy or Milk protein ingestion in humans. *J Nutr.* 2003; 133:1308–15.
17. Atherton PJ, Smith K, Etheridge T, Rankin D, Rennie MJ. Distinct anabolic signalling responses to amino acids in C2C12 skeletal muscle cells. *Amino Acids.* 2010;38:1533–9.
18. Witard OC, Jackman SR, Breen L, Smith K, Selby A, Tipton KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr.* 2014;99:86–95.

-
19. Fogelholm M. Dairy products, meat and sports performance. *Sports Med.* 2003;33:615–31.
20. Hoffman JR, Falvo MJ. Protein - Which is Best? *J Sports Sci Med.* 2004;3:118–30.
21. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ.* 2009;339:b2535.

Page

22. National Library of Medicine. Medical Subject Headings. MeSH Descriptor Data. 2018 [cited 2018 June 12]. Available from: <https://www.ncbi.nlm.nih.gov/mesh/?term=dairy+product>